

Eurasian Watermilfoil

Myriophyllum spicatum

Eurasian watermilfoil is a non-native aquatic plant found in much of the shallow areas of most lakes in the region. It is suspected this incredibly invasive aquatic weed was recently introduced into nearby lakes from other parts of the Columbia River drainage, most likely on boat trailers. Because it is widely distributed and difficult to control, Eurasian watermilfoil is considered to be the most serious aquatic weed problem in the Northwest.

As its name suggests it has its origins in Europe and Asia. It probably arrived in the United States in the 1940s and has quickly spread to all but a few states. Eurasian watermilfoil, commonly referred to as simply watermilfoil, (there is also a less invasive native watermilfoil), is actually an attractive plant that was once commonly sold as an aquarium plant.

Even a tiny fragment of the plant is enough to establish a colony that can quickly out compete beneficial native aquatic plants. These massive rooted colonies can become so dense that they not only compromise the water quality, but can suffocate fish and destroy spawning habitat for some species.

Watermilfoil also has a serious impact on an areas economic well being. Most water based recreational activities are affected by it to some degree. Boats can have trouble navigating through the nearly solid floating mats occasionally getting stuck. Fish populations can decline and swimmers have been known to drown in rare instances. Dense floating mats can also clog the inlets of power generating plants and increase the cost of maintenance.

One of the leading causes of the proliferation of Eurasian watermilfoil is excessive nutrient input into the lake and it's tributaries both from improper land use practices and from individual home owners. As you read through the Lake*A*Syst materials keep in mind the effect of nutrient runoff, especially nitrogen and phosphorous, and the effect on the water quality of Hauser Lake. What you do or don't do can make a difference.

Control of Eurasian Watermilfoil

Once watermilfoil becomes well established within a body of water, it is difficult or impossible to remove. There has been some success using aquatic herbicides, however significant control is difficult considering the expense of herbicide application.

Timing the application is also critical. Keep in mind that the application by a private landowner of any herbicide to a body of public water or a tributary of public water is **illegal** no matter what the herbicide package may say.

One method of helping prevent the spread of milfoil is utilizing the **Hauser Lake Watercraft Rinse Station**. Lead by local citizens of the Hauser Lake Watershed Coalition, Idaho's first Rinse Station was built in May of 2007. When properly utilized by boaters, this facility will help prevent the infestation of many kinds of invasive species. Do your part to protect your lake...**Rinse Before You Launch**.

Identification

As a group, watermilfoils are easy to identify, however identifying the exact species is more challenging. All watermilfoil have feather-like leaves arranged in a whorl of four leaves around the stem. Here are some tips to identify Eurasian Watermilfoil from the native milfoils. The Eurasian variety usually has twelve or more leaflets per leaf while the native northern variety has fewer than ten. The leaves of the Eurasian variety tend to collapse around the stem while the northern species tend to remain more rigid when removed from the water. Also the stem is generally more reddish with the Eurasian species.

